



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

VII. *A finite and exact expression for the Refraction of an Atmosphere nearly resembling that of the Earth.* By THOMAS YOUNG, M. D. For. Sec. R. S.

Read February 5, 1824.

IT has lately been demonstrated, in the Journal of the Royal Institution, that if the pressure of the atmosphere,  $y$ , be represented either by the square or by the cube of the square root of the density,  $z$ , the astronomical refraction,  $r$ , may be obtained in a finite equation. Mr. IVORY, in a very ingenious and elaborate paper lately presented to the Royal Society, has computed the refraction, by means of several refined transformations, and with the assistance of converging series, from an equation which expresses the pressure in terms of the density and of its square: I have now to observe, that if we substitute, for the simple density, the cube of its square root, and make  $y = \frac{3}{2} z^{\frac{3}{2}} - \frac{1}{2} z^2$ , we shall represent the constitution of the most important part of the atmosphere with equal accuracy, although this expression supposes the total height somewhat smaller than the truth, and belongs to one of those hypotheses, which Mr. IVORY has considered as inadmissible: it has the advantage, however, of affording a direct equation for the refraction, which agrees very nearly with Mr. IVORY's table, and still more accurately with the French table, and with that which has been published for some years in the Nautical Almanac.

Since  $dx = \frac{-dy}{mz}$ ,  $m$  being the number of times that the modulus of the atmospherical elasticity is contained in the

160 *Dr. YOUNG on a finite and exact expression for the refraction*

radius of the earth, and here  $dy = \frac{9}{4} \sqrt{z} dz - z dz$ , we have  $dx = -\frac{9}{4} \frac{dz}{m \sqrt{z}} + \frac{dz}{m}$ , and  $\int dx = -\frac{9}{2m} \sqrt{z} + \frac{z}{m} + \frac{7}{2m}$ , for the height above the earth's surface, which, when  $z = 0$ , becomes  $\frac{7}{2} \times 27300 = 95550$  feet. For the refraction, we have the equation  $dr = \frac{-p dz}{\sqrt{(x^2 - s^2 - 2p[1-z])}}$  (Astr. Coll. XV.)  $= \frac{-p dz}{\sqrt{(2f dx + v^2 - 2p + 2pz)}}$ , which is the value originally assigned to this fluxion by Dr. BROOK TAYLOR;  $v$  being the sine of the apparent altitude; and here

$$dr = \frac{-p dz}{\sqrt{\left(\frac{7}{m} - \frac{9}{m} \sqrt{z} + \frac{2z}{m} + v^2 - 2p + 2pz\right)}}$$

or, if  $\sqrt{z} = \psi$ , and  $dz = 2\psi d\psi$ ,

$$-\frac{dr}{2p} = \frac{\psi d\psi}{\sqrt{\left(\frac{7}{m} - 2p + v^2 - \frac{9}{m} \psi + \left[\frac{2}{m} + 2p\right] \psi^2\right)}}$$

which is equivalent to the  $\frac{x dx}{\sqrt{(a + bx + cx^2)}}$  of the Article FLUENTS in the *Encyclopædia Britannica*, No. 259; the fluent being

$$\frac{1}{c} \left[ \sqrt{(a + bx + cx^2)} - \frac{b}{2\sqrt{c}} \text{hl}(2cx + b + 2\sqrt{c}\sqrt{a + bx + cx^2}) \right]$$

and its whole value, from  $z = 1$  to  $z = 0$ , being  $-\frac{cr}{2p}$

$$= \sqrt{(a' + v^2)} - v + \frac{b}{2\sqrt{c}} \text{hl} \frac{2c + b + 2v\sqrt{c}}{b + 2\sqrt{c}\sqrt{(a' + v^2)}}, \text{ putting } a' = \frac{7}{m} - 2p, \text{ since } a + b + c = v^2.$$

For the numerical values of the coefficients, taking, at the temperature of  $50^\circ$ ,  $p = .0002835$ , and  $\frac{1}{m} = .001294 = \frac{1}{772.8}$ ,

$$a = \frac{7}{m} - 2p + v^2 = .008491 + v^2, b = \frac{-9}{m} = -.011646,$$

$$\text{and } c = \frac{2}{m} + 2p = .003155; \text{ hence } \frac{2p}{c} = .17972, \sqrt{a'} = .0921466, \sqrt{c} = .05617, \frac{-b}{2\sqrt{c}} = .10367, 2c + b = -.005336, \text{ and}$$

$$r = .17972 \left( \sqrt{(.008491 + v^2)} - v - .10367 \text{hl} \frac{.005336 - .11234v}{.011646 - .11234\sqrt{(.008491 + v^2)}} \right);$$

and at the horizon, when  $v = 0$ ,

$$r = .17972 \left( .0921466 - .10367 h l \frac{.005336}{.011646 - .11234 \times .092147} \right) =$$

$.009840 = 33' 42'',5$ ; which is only  $1'',5$  less than the quantity assigned by the French tables and in the Nautical Almanac, while Mr. IVORY makes it  $34' 17'',5$ . Again, if we take  $v = .1$ , for the altitude  $5^\circ 44' 21''$ , we obtain  $8' 49'',5$  for the refraction, while the Nautical Almanac gives us  $8' 53''$ , and Mr. IVORY's table  $8' 49'',6$ . There is however no reason for proceeding to compute a new table by this formula, the method employed for the table in the Nautical Almanac being rather more compendious in all common cases: and even if it were desired to represent Mr. IVORY's table by the approximation there employed, we might obtain the same results, with an error never much exceeding a single second, from the equation

$$00028333 = \frac{v}{s} r + \frac{2.26 + \frac{1}{2}vv}{ss} r^2 + 5400 \frac{rr}{ss} \left( \frac{v}{s} r + \frac{1.13 + \frac{1}{2}vv}{ss} r^2 \right).$$

*Welbeck Street,*

*3rd. February, 1824.*